

**NPAFC
Doc. 442
Rev.**

**INCIDENTAL CATCHES OF SALMONIDS BY U.S. GROUND FISH FISHERIES
IN THE BERING SEA/ALEUTIAN ISLANDS, GULF OF ALASKA,
AND THE PACIFIC COAST, 1990-1999**

by

Jerald D. Berger
Resource Ecology and Fisheries Management Division
Alaska Fisheries Science Center
National Marine Fisheries Service

Submitted to the

NORTH PACIFIC ANADROMOUS FISH COMMISSION

by the

United States Party

October 1999

THIS PAPER MAY BE CITED IN THE FOLLOWING MANNER:

Berger, J.D. 1999. Incidental catches of salmonids in U.S. groundfish fisheries in the Bering Sea/Aleutian Islands, Gulf of Alaska, and Pacific Coast Areas, 1990-1999. (Doc. 442) 6p. Resource Ecology and Fisheries Management Div., Alaska Fish. Sci. Cent., NMFS, NOAA, U.S. Dep. Commerce, 7600 Sand Point Way NE, Seattle, WA 98115-0070.

**INCIDENTAL CATCHES OF SALMONIDS BY U.S. GROUND FISH FISHERIES
IN THE BERING SEA/ALEUTIAN ISLANDS, GULF OF ALASKA,
AND THE PACIFIC COAST, 1990-1999**

ABSTRACT

Incidental catches of Pacific salmonids in U.S. groundfish trawl fisheries of Alaska and the Pacific West Coast are presented for 1977 through October 9, 1999. Estimated numbers of salmonids caught incidentally in the U.S. groundfish fisheries in the Bering Sea/Aleutian Islands were 30,501 in 1990; 79,133 in 1991; 83,405 in 1992; 289,210 in 1993; 140,811 in 1994; 44,859 in 1995; 141,200 in 1996; 117,753 in 1997; and 128,203 in 1998. In the Gulf of Alaska, incidental catches were 21,085 in 1990; 53,844 in 1991; 38,308 in 1992; 80,853 in 1993; 54,486 in 1994; 79,439 in 1995; 19,937 in 1996; 18,539 in 1997; and 30,480 in 1998. The Pacific Coast U.S. groundfish fisheries off Washington, Oregon, and California caught 617 salmon in 1990; 6,331 salmon in 1991; 5,100 salmon in 1992; 8,373 salmon in 1993; 4,001 salmon in 1994; 15,991 in 1995; 3,437 in 1996; and 5,410 in 1997. Through October 9, 1999, the incidental catches were 79,233 salmon in the Bering Sea/Aleutian Islands and 32,236 salmon in the Gulf of Alaska. The estimates of salmon caught off the Pacific Coast in 1998 and 1999 are currently unavailable. The last joint venture operations took place in 1990 in the Bering Sea/ Aleutian Islands and off the Pacific Coast. Their incidental catches of salmon were 152 and 9,308, respectively.

INTRODUCTION

Groundfish fisheries are conducted primarily by the use of trawl gear that occasionally encounters and catches Pacific salmonids. Other gear used to harvest groundfish - longline and pots - generally do not catch many salmonids. U.S. fishing regulations consider salmonids to be prohibited species for the groundfish fisheries, and when taken, they must immediately be returned to the sea. The incidental catches of salmonids are monitored by the U.S. National Marine Fisheries Service (NMFS) and estimated by its scientific observer program. At present, there are no specific limits set for salmonid bycatch in the Gulf of Alaska nor off the Pacific Coast. In the Bering Sea, a catch of 42,000 non-Chinook salmon during ? B Season? triggers a time-area closure. This was established in 1994 to minimize the incidental catches of chum salmon (*Onchorhynchus keta*). Additionally, incidental catch rates of individual vessels and the fleet average are monitored closely and made known to the fleet, so individual vessels can take voluntary actions to minimize their bycatch.

The observer program coverage for the groundfish fisheries off Alaska is 100% for the large-sized vessel fleet (greater than 125 ft. length overall) and 30% for the medium-sized vessel fleet (60-125 ft. length overall). The small-sized vessel fleet (less than 60 ft. length overall) is not monitored by direct observer coverage. Instead, some of their catches are sampled when they deliver to motherships or shore-based processing plants. Motherships are staffed 100% of the time by observers and shore-based plants are staffed 30%-100%, depending upon the volume of fish processed. For the Pacific Coast, only the at-sea processors (catcher-processors and motherships) are included in these estimates.

RESULTS

Observers count salmonids, take length-weight measurements of salmonid samples, and estimate total groundfish catch weight. Tables 1-4 summarize the estimates from the observer sampling.

Table 1. Total groundfish catch and estimated bycatch of chinook and other Pacific salmon in U.S. groundfish fisheries, 1990-1999*.

Year	Region	Groundfish Catch (mt)	Chinook -----	Chum -----	Coho -----	Sockeye -----	Pink -----	Total -----
Domestic								
1990	BSAI	1,706,379	14,085	16,202	153	30	31	30,501
1991	BSAI**	2,154,903	48,873	29,706	396	79	79	79,133
1992	BSAI	1,963,523	41,955	40,090	1,266	14	80	83,405
1993	BSAI**	1,754,384	45,964	242,895	321	22	8	289,210
1994	BSAI	1,855,031	44,380	95,978	231	20	202	140,811
1995	BSAI	1,830,295	23,079	20,901	858	0	21	44,859
1996	BSAI	1,755,872	63,205	77,771	218	5	1	141,200
1997	BSAI	1,740,663	50,218	67,349	114	3	69	117,753
1998	BSAI	1,531,838	58,966					-----69,237---
-----		128,203						
1999	BSAI	1,243,051	16,861					-----62,372---
-----		79,233						
1990	GOA	244,397	16,913	2,541	1,482	85	64	21,085
1991	GOA	269,616	38,894	13,713	1,129	51	57	53,844
1992	GOA	269,797	20,462	17,727	86	33	0	38,308
1993	GOA	255,434	24,465	55,268	306	15	799	80,853
1994	GOA	239,503	13,973	40,033	46	103	331	54,486
1995	GOA	216,585	14,647	64,067	668	41	16	79,439
1996	GOA	202,054	15,761	3,969	194	2	11	19,937
1997	GOA	230,448	15,119	3,349	41	7	23	18,539
1998	GOA	245,115	16,941					-----13,539---
-----		30,480						
1999	GOA	217,680	24,943					-----7,293---
-----		32,230						
1990	WOC***	4,478	617	0	0	0	0	617
1991	WOC***	198,953	6,165	11	132	0	23	6,331
1992	WOC***	155,333	4,863	36	201	0	0	5,100
1993	WOC***	99,698	4,843	71	20	126	3,313	8,373
1994	WOC***	175,731	3,626	275	39	0	61	4,001
1995	WOC***	103,598	11,577	200	1,775	6	2,433	15,991
1996	WOC***	129,489	3,152	196	89	0	0	3,437
1997	WOC***	147,221	4,404	139	304	0	563	5,410
1998	WOC***	147,396	na	na	na	na	na	na
1999	WOC***	140,001	na	na	na	na	na	na
Joint Venture								
1990	BSAI	133,438	147	2	3	0	0	152
1990	GOA	0	0	0	0	0	0	0
1990	WOC	172,069	9,141	133	34	0	0	9,308

Regions: BSAI = Bering Sea/Aleutian Islands
 GOA = Gulf of Alaska
 WOC = Washington-Oregon-California

* Through 10/9/99

** 1 steelhead was reported in 1991 and 5 steelheads were reported in 1993.

*** At sea processing only

na - Not available

Note: For 1998 and 1999, the estimates of non-chinook salmon have not yet been separated by species and are thus listed as a single value.

Table 2. Average body weight of chinook salmon incidentally caught in U.S. groundfish fisheries, 1990-1999*.

Year	Region	Male Average Weight (kg)	Female Average Weight (kg)
1990	BSAI	3.74	3.87
1991	BSAI	2.83	3.03
1992	BSAI	3.26	3.32
1993	BSAI	3.76	4.07
1994	BSAI	4.37	4.29
1995	BSAI	4.95	3.97
1996	BSAI	3.68	3.90
1997	BSAI	4.00	4.16
1998	BSAI	2.91	3.30
1999	BSAI	3.43	3.83
1990	GOA	3.57	3.88
1991	GOA	3.03	3.43
1992	GOA	3.46	3.32
1993	GOA	2.84	2.77
1994	GOA	3.70	3.68
1995	GOA	4.48	3.31
1996	GOA	2.71	2.96
1997	GOA	4.28	4.10
1998	GOA	3.00	3.07
1999	GOA	2.45	3.12
1990	WOC	2.14	2.65
1991	WOC	3.18	3.40
1992	WOC	2.93	4.08
1993	WOC	2.22	2.88
1994	WOC	2.26	2.69
1995	WOC	2.72	2.64
1996	WOC	4.67	5.35
1997	WOC	2.71	2.92
1998	WOC	3.06	3.68
1999	WOC	3.48	2.81

Regions: BSAI = Bering Sea/Aleutian Islands

GOA = Gulf of Alaska

WOC = Washington-Oregon-California

* Data presented for 1999 are based on preliminary data.

Table 3. Average body weight of chum salmon incidentally caught in U.S. groundfish fisheries, 1990-1999*.

Year	Region	Male Average weight (kg)	Female Average weight (kg)
1990	BSAI	2.28	2.18
1991	BSAI	2.44	2.29
1992	BSAI	2.53	2.51
1993	BSAI	2.16	1.96
1994	BSAI	2.72	2.65
1995	BSAI	3.45	3.17
1996	BSAI	2.47	2.24
1997	BSAI	2.35	2.24
1998	BSAI	2.89	2.50
1999	BSAI	2.89	2.91
1990	GOA	3.36	3.11
1991	GOA	3.11	2.89
1992	GOA	3.32	3.06
1993	GOA	2.40	2.27
1994	GOA	3.40	3.20
1995	GOA	3.14	3.12
1996	GOA	3.09	2.43
1997	GOA	5.12	4.01
1998	GOA	3.84	3.77
1999	GOA	4.24	4.12
1990	WOC	--**	--**
1991	WOC	4.55	4.35
1992	WOC	4.58	4.55
1993	WOC	3.37	1.07***
1994	WOC	5.35	5.27
1995	WOC	1.80	2.95
1996	WOC	5.05	4.22
1997	WOC	3.12	3.18
1998	WOC	4.15	3.77
1999	WOC	5.74	4.77

Regions: BSAI = Bering Sea/Aleutian Islands
 GOA = Gulf of Alaska
 WOC = Washington-Oregon-California

* Data presented for 1999 are based on preliminary data.
 ** None caught.
 *** Very few fish, mostly very small.

Table 4. Estimated incidental catches of Pacific salmonids previously reported to the International North Pacific Fisheries Commission, 1977-1989.

Year	Bering Sea/ Aleutian*	Gulf of Alaska**	Washington- Oregon-California*
1977	47,840	5,222	14,627
1978	44,548	45,603	5,924
1979	107,706	21,460	8,666
1980	122,002	36,069	8,433
1981	43,191	30,860	11,474
1982	23,623	6,967	11,798
1983	42,666	13,874	5,143
1984	84,138	75,846	10,255
1985	20,423	14,102	2,288
1986	20,983	20,820	43,790
1987	14,234	1,221	13,285
1988	9,380	147	16,168
1989	14,153	0	9,199

* Reported in INPFC Doc. 3588, October 1990

** Reported in INPFC Doc. 3417, September 1989