

Impact of Predation by Salmon Sharks (*Lamna ditropis*) and Daggertooth (*Anotoapterus nikparini*) on Pacific Salmon (*Oncorhynchus* spp.) Stocks in the North Pacific Ocean

Kazuya Nagasawa¹, Tomonori Azumaya², and Yukimasa Ishida³

¹ Nikko Branch, National Research Institute of Aquaculture, Fisheries Research Agency, Nikko, Tochigi 321-1661, Japan

² National Hokkaido Fisheries Research Institute, Fisheries Research Agency, Kushiro, Hokkaido 085-0802, Japan

³ Kuroshio Research Laboratory, National Research Institute of Fisheries Science, Fisheries Research Agency, Kochi 780-8010, Japan



Keywords: Predation impact, salmon shark, daggertooth, salmon stocks, North Pacific.

Various species of fishes, seabirds and marine mammals are known as predators of Pacific salmon (*Oncorhynchus* spp.) in the North Pacific Ocean. However, there is little quantitative information about impact of predation by such predators on Pacific salmon populations. In this paper, we focus on predation by salmon sharks (*Lamna ditropis*) in oceanic offshore waters of the North Pacific Ocean and report on possible substantial impact of their predation on salmon stocks, based on the recent increase in salmon shark abundance on the high seas of the North Pacific Ocean. Also, we discuss predation impact by daggertooth (*Anotoapterus nikparini*, formerly *A. pharao*) on salmon stocks, based on current information on their abundance and frequency of occurrence of slash marks on the lateral side of high-seas salmon caused by the attack of daggertooth.

We calculated annual CPUE (number/1000 tan of surface gillnets) of salmon sharks using the data collected by Japanese salmon research vessels that operated in the western and central North Pacific Ocean from 1972 to 2000. Salmon shark CPUE was low from 1984 to 1993 but increased sharply in 1996 and thereafter remained at a high level. A similar increase in salmon shark abundance has been reported from the Gulf of Alaska (Wright and Hulbert 2000). These results indicate that the North Pacific salmon shark population has recovered since the mid- or late 1990s. We estimate that the loss of high-seas salmon due to salmon shark predation was approximately $73\text{-}146 \times 10^6$ salmon ($113\text{-}226 \times 10^3$ metric tons) in 1989 when salmon shark CPUE was low (Nagasawa 1998). Thus, there is a possibility that the current increase in salmon shark abundance in the whole region of the North Pacific has increased mortality of salmon during the high-seas phase of the life history.

Daggertooth is a species that has been recently noticed as a predator of offshore salmon (Welch et al. 1991; Radchenko and Semenchenko 1996). Current Russian surface trawl surveys indicate that the abundance of daggertooth is not low in the western North Pacific and that it feeds on large numbers of salmon (Melnikov 1997). Slash marks from daggertooths are not frequently but sometimes found on salmon from high seas and coastal waters. The incidence of slash marks on salmon is usually low, ranging from less than 1% to 4 or 5%, in offshore waters (Ishida et al. 1991) but slightly higher (up to 12%) in coastal waters (Gilhousen 1989; Henderson et al. 1990). The species is widely distributed in offshore waters of the North Pacific Ocean (Nagasawa 1992), and its abundance seems to have been increasing since the early 1990's. Based on this information, we can regard daggertooth as significant salmon predators as well.

REFERENCES

- Gilhousen, P. 1989. Wounds, scars and marks on Fraser River sockeye salmon with some relationships to predation losses. Int. Pac. Salmon Fish. Comm. Rep. 42. 64 p.
- Ishida, Y., K. Tadokoro, and K. Nagasawa. 1991. Preliminary estimates of the incidence of slash marks on salmonids caught by Japanese research vessels in the North Pacific Ocean and Bering Sea in 1991. (Document submitted to the Annual Meeting of the International North Pacific Fisheries Commission, Tokyo, October 1991). National Research Institute of Far Seas Fisheries, Shimizu. 3 p.
- Henderson, M., S. McKinnell, and L. Margolis. 1990. Preliminary estimates of the incidence of slash marks on sockeye salmon (*Oncorhynchus nerka*) caught in commercial fisheries in British Columbia in 1990. (Document submitted to the Annual Meeting of the International North Pacific Fisheries Commission, Vancouver, November 1990. Department of Fisheries and Oceans, Pacific Biological Station, Nanaimo. 8 p.

- Melnikov, I.V. 1997. Pelagic fishes-predator as consumers of Pacific salmon: distribution in Russian economic zone and adjacent waters: abundance and some biological features. *Izv. TINRO* 122: 213–233. (In Russian with English abstract).
- Nagasawa, K. 1992. Distribution of daggertooth *Anotopterus pharao* in the North Pacific Ocean. *Salmon Rep. Ser.* 36: 93–99. (In Japanese and English) [Available from the National Research Institute of the Far Seas Fisheries, Shimizu-shi, shizuoka, 424-9633, Japan]
- Nagasawa, K. 1998. Predation by salmon sharks (*Lamna ditropis*) on Pacific salmon (*Oncorhynchus* spp.) in the North Pacific Ocean. *N. Pac. Anadr. Fish Comm. Bull. No. 2*: 419–433.
- Radchenko, V.I., and A.Yu. Semenchenko. 1996. Predation of daggertooth on immature Pacific salmon. *J. Fish Biol.* 49: 1323–1325.
- Welch, D.W., L. Margolis, M.A. Henderson, and S. McKinnell. 1991. Evidence for attacks by the bathypelagic fish *Anotopterus pharao* (Myctophiformes) on Pacific salmon (*Oncorhynchus* spp.). *Can. J. Fish. Aquat. Sci.* 48: 2403–2407.
- Wright, R.A., and L. Hulbert. 2000. Shark abundance increases in the Gulf of Alaska. *PICES Press*, 8(2): 16–17, 22.